

Restoring Nature's Song: How Habitat Restoration Benefits Avifauna

Sriyani Wickramasinghe*, B.A.D.D.P Bopearachchi

Department of Biological Science, Faculty of Applied Sciences, Rajarata University of Sri Lanka, Mihintale

Abstract

Habitat restoration has proved to be a vital strategy in combating the widespread biodiversity loss caused by habitat degradation across the globe. The key roles played by avian species in critical ecosystem functions such as pollination, seed dispersal, and decomposition have made them an integral part of ecosystems, making them vital to the restoration process. Their detectability and cost-effective survey methods have positioned avifauna as focal species in ecological restoration research. This article delves into the effects of habitat restoration on avifaunal distribution, revealing a tapestry of positive outcomes. Restored habitats witness increased avian community diversity and evenness, heightened habitat heterogeneity, and greater densities of avian and migratory species. Avifauna's presence also facilitates breeding success, reduces nest parasitism, and fosters additional foraging and breeding grounds. Furthermore, habitat restoration has shown remarkable successes in reviving songbird populations in fragmented forests and rescuing endangered species like red-crowned cranes. In light of the transformative impact of avian restoration, understanding the effects of restoration on avifauna is pivotal in guiding management activities of restored sites. Implementing strategic management approaches such as predator control, maintaining habitat heterogeneity, controlling nest parasites, reintroducing natural disturbance regimes, and selecting appropriate plant species ensures a nurturing environment for avian survival. Additionally, promoting a heterogeneous environment in aquatic systems has proved crucial in enhancing bird diversity. Improving ongoing research and adaptive management practices further empowers restoration efforts, allowing us to compose a symphony of avian abundance, resilience, and harmonious coexistence with the natural world.

Keywords: Habitat Restoration, Avifauna, Biodiversity, Ecosystem Services, Ecological Resilience

1. Introduction

The rhythmic symphony of nature's song has been gradually silenced by the rapid degradation of habitats worldwide. Land clearing and environmental degradation driven by human activities have resulted in the loss of biodiversity, threatening the delicate balance of ecosystems (Macias, 2009; Ortega-Álvarez and Lindig-Cisneros, 2012). In response to this ecological crisis, habitat restoration has emerged as a beacon of hope, offering a chance to reverse the trajectory of biodiversity loss.

At the heart of restoration efforts, avifauna stands as a focal point in the quest to reclaim our natural heritage. Over the last few decades, ecologists have increasingly recognized the indispensable role of avian species in restoring ecosystems (Majer, 2009; Alsila *et al*, 2020). Birds, with their unique abilities

Correspondence: sriwick@gmail.com

ISSN 2235-9362 Online ©2023 University of Sri Jayewardenepura

to influence critical ecological processes such as pollination, seed dispersal, and decomposition, become the vital threads that interweave the fabric of restored habitats.

The significance of avifauna in ecological restoration research extends beyond their ecological functions. These feathered creatures, with their vocal presence and distinctive attributes, become valuable indicators of restoration success. Their detectability and cost-effective survey methods enable researchers to assess the impact of restoration efforts on avian populations over vast areas (Gardali *et al.*, 2006). Additionally, the wide range of ecological requirements exhibited by different avian species offers a window into the overall health and functionality of restored ecosystems (Gaines *et al.*, 2007; Johnson and Sandercock, 2010).

In this article, we embark on a journey to explore the profound effects of habitat restoration on avifaunal distribution. We delve into the successes and challenges faced in avian restoration, seeking to understand how these tireless guardians of our ecosystems respond to the revival of their habitats. Moreover, we unravel the web of management activities that guide restored sites towards becoming thriving sanctuaries for avian biodiversity.

Our goal is to emphasize on the vital role played by avifauna in the restoration process, and show how their presence is crucial in shaping the future of our planet's ecosystems. By appreciating their contributions to key ecosystem functions (pollination, seed dispersal, and decomposition), we tend to broaden the understanding of intricate connections that forms a functioning ecosystem.

As we venture through the pages that follow, we call upon all to join hands in this collective endeavor. By understanding the significance of habitat restoration in preserving avian biodiversity, we can take actions that reverberate far beyond the horizon of our vision. Let us, as stewards of this planet, rise together to restore nature's song and ensure the harmonious coexistence of avifauna and humanity.

2. Methodology

The primary objective of this literature review is to explore the effects of habitat restoration on avifaunal distribution and its significance in restoring ecosystems. The research question guiding this review is: "How do habitat restoration impact avifauna populations and ecosystem functions?" A comprehensive search was done in an array of academic databases, including PubMed, Scopus, Web of Science, and Google Scholar. (Keywords used -"habitat restoration," "avifauna," "bird populations," "ecosystem functions," and related terms). The search was limited to articles published in English from the last two decades (1997 to 2021) to ensure relevance to current practices and research. Data were extracted from the selected sources using a systematic approach. Key findings, trends, and outcomes regarding avian responses to habitat restoration were recorded. Information on avifaunal diversity, population dynamics, breeding success, and ecological functions (pollination, seed dispersal, decomposition) were synthesized. The collected data were analyzed using thematic analysis. Similarities and differences in avian responses to habitat restoration were identified. Themes related to the positive impact of restoration on avifauna, challenges faced, and successful case studies were organized for synthesis. All information and data used in this literature review are properly cited and credited to the original authors and sources. The review might not cover every relevant study on the topic due to the vastness of the literature available.

3. Results and Discussion

3.1 *The Role of Avifauna in Restoration Ecology*

Avifauna, or bird species, emerges as a keystone element in the success of habitat restoration efforts, playing a pivotal role in restoring critical ecosystem functions. As ecosystem engineers, birds exert substantial influence over key ecological processes, including pollination, herbivory, seed dispersal, and decomposition.

One of the essential roles avifauna play in restoration ecology is pollination. Many bird species, such as hummingbirds and sunbirds, act as efficient pollinators, transferring pollen between flowers as they forage for nectar (Balvanera, 2005; Sekercioglu, 2006). Through this process, they facilitate plant reproduction, enhancing genetic diversity within restored habitats and promoting the growth of diverse plant communities.

Herbivory is another ecological process influenced by avian species. Birds, particularly granivores and frugivores, consume seeds and fruits from various plant species (Coley and Barone, 1996). In doing so, they shape plant populations, control seed dispersal, and influence vegetation dynamics within restored ecosystems. By curbing the growth of certain plant species and promoting the proliferation of others, avifauna contribute to the balanced structure and composition of plant communities.

Seed dispersal is a vital service provided by birds, allowing plants to colonize new areas and expand their range. Avian species, especially those with specialized adaptations for seed dispersal, disperse seeds over large distances, aiding in the establishment of new plant populations (Wunderle, 1997; Bollen, 2004). In the context of restoration, this process becomes paramount in accelerating the colonization of restored habitats by native plant species, facilitating the recovery of biodiversity.

Furthermore, avifauna significantly contributes to the process of decomposition. Birds, particularly scavengers, play an important role in nutrient recycling by feeding on carrion which, aids in the breakdown of organic matter (Chapman *et al.*, 2003; Botes *et al.*, 2006). This accelerates the release of nutrients back into the ecosystem, enriching the soil which, supports the growth of vegetation within restored habitats.

Given the critical ecological functions that they perform, avifauna is considered a focal species in ecological restoration research. Their distinctive attributes make them well-suited subjects for monitoring and research within restored ecosystems. Detecting and identifying bird species is relatively straightforward due to their vocalizations, facilitating their inclusion in post-restoration assessments (Gardali *et al.*, 2006). Additionally, avifauna can be surveyed cost-effectively over large areas, providing valuable data on biodiversity and habitat quality across restoration sites.

Moreover, avian species exhibit a wide range of ecological requirements, occupying diverse habitats and responding to changes in ecosystem structure and composition (Gaines *et al.*, 2007; Johnson and Sandercock, 2010). This diversity allows researchers to explore a variety of restoration scenarios and gauge the response of different avian species to specific restoration strategies. By studying avifauna within restoration efforts, ecologists gain valuable insights into the overall health and functionality of restored ecosystems.

3.2 *The Value of Management Activities in Restored Sites*

The success of habitat restoration in benefiting avifauna is closely intertwined with effective management activities. Understanding the effects of restoration on avifauna becomes the compass that guides these vital interventions, ensuring that restored sites become thriving sanctuaries for our feathered friends.

One of the critical roles of understanding the effects of restoration on avifauna lies in tailoring management strategies to meet the specific needs of bird species. Avian responses to restoration efforts are species-specific (Gaines *et al.*, 2007; Johnson and Sandercock, 2010; Alsila *et al.*, 2020), and understanding these nuances allows us to implement targeted approaches that maximize positive outcomes. By taking into account the diverse ecological requirements and behaviors of different avian species, management activities can be fine-tuned to create a supportive environment for their survival and growth.

A range of management strategies have proven to be valuable in nurturing avifauna within restored habitats. Predator control emerges as a key approach to promote adult bird survival and reduce nest predation. Implementing predator control projects and enhancing habitat quality at the landscape level can help protect nesting sites and safeguard vulnerable bird populations (Fletcher *et al.*, 2006; Fletcher and Koford, 2003).

Maintaining habitat heterogeneity stands as another vital aspect of avian restoration. Restored ecosystems benefit from a mosaic of diverse habitats, each providing unique resources and niches for different bird species. By preserving a variety of vegetation structures, such as grasslands, shrub lands, and forests, avifauna find ample foraging and nesting opportunities, contributing to increased avian diversity and overall ecosystem resilience (Seigel *et al.*, 2005; Ortega-Álvarez *et al.*, 2013).

Controlling nest parasites is an essential management activity that reduces the pressure of parasitism on avian populations. By managing and monitoring avian nest parasite species, we can help mitigate the negative impact of parasitism on nesting success (Small *et al.*, 2007). This, in turn, supports healthier avian communities within restored habitats.

Reintroducing natural disturbance regimes is a crucial step in maintaining habitat heterogeneity and promoting avian diversity. Many bird species have evolved alongside natural disturbances, such as fires or floods, and depend on these events for nesting opportunities and foraging grounds (Masters *et al.*, 2000; Brawn, 2006). By reintroducing such natural processes in controlled settings, we can mimic the dynamic nature of undisturbed ecosystems and foster thriving avifauna populations.

The choice of plant species during the early stages of restoration significantly impacts avian colonization. Selecting fast-growing successional plant species that provide food resources and suitable habitats can accelerate the arrival of bird species to restored sites (Twedt *et al.*, 2002; Flesch and Esquer, 2020). These pioneer species act as beacons, attracting avifauna to the newly restored habitats and facilitating the establishment of diverse bird communities.

For aquatic systems, promoting a heterogeneous environment becomes particularly crucial in enhancing bird diversity. A mix of vegetation types, prey availability, and water depths in aquatic habitats fosters a rich habitat mosaic that appeals to a wide range of water bird species (Fink *et al.*, 2009). By creating such varied conditions, avian diversity thrives, and wetland sanctuaries become abundant hubs of life and activity.

3.3 Success Stories of Avian Restoration

Habitat restoration efforts have yielded remarkable success stories, demonstrating the potential for positive impacts on avifauna and the restoration of endangered bird populations. Among these triumphs, two standout examples showcase the resilience of avian species in the face of adversity: the rescue of red-crowned cranes and the revival of songbirds in fragmented forests.

Rescue of Red-Crowned Cranes:

The red-crowned crane (*Grus japonensis*) stands as an iconic symbol of conservation success in East Asia. Once on the brink of extinction due to habitat loss and hunting, this majestic bird faced dire prospects for survival. However, dedicated restoration efforts have turned the tide in its favor.

In China, coastal wetlands at Yancheng, and Tsurui Ito Tancho Sanctuary in Hokkaido has played a pivotal role in the rescue of red-crowned cranes. These areas provide vital wetland habitats and abundant food resources, offering a secure breeding ground for these magnificent birds (Higuchi *et al.*, 1998 Momose and Momose, 2019). Through targeted management activities, including habitat restoration, predator control, and community involvement, the red-crowned crane population has rebounded significantly, bringing hope for the species' future.

Revival of Songbirds in Fragmented Forests:

Fragmentation of forests due to human activities has been a significant threat to numerous songbird species worldwide. Isolated forest patches often lead to reduced genetic diversity and increased vulnerability to predation, posing severe challenges to bird populations. Nevertheless, restoration initiatives that focus on reconnecting fragmented habitats have sparked incredible success stories. In Brazil, the Atlantic Forest has been a focal point for such efforts. The restoration of degraded areas and the establishment of biological corridors have facilitated the movement of songbirds between forest fragments (Ribeiro *et al.*, 2009). This connectivity has not only enhanced genetic diversity but also allowed for the re-establishment of populations in once-isolated patches. Species like the golden-capped parakeet (*Aratinga auricapilla*) and the black-fronted piping-guan (*Pipile jacutinga*) have shown encouraging signs of recovery in restored areas (Ribeiro *et al.*, 2009).

Among the avian species observed, the sighting of the Cheer Pheasant (*Catreus wallichii*), a species endemic to the Western Himalayan Endemic Bird Area, holds special significance. This sighting underscores the value of the restored forest system, as the Cheer Pheasant is classified as Vulnerable on the IUCN Red List due to threats like habitat loss and illegal hunting. Moreover, other endemic species like the Yellow-breasted Greenfinch, Scarlet Finch, Brown-fronted Woodpecker, Slaty-headed Parakeet, and Himalayan Bulbul were also documented during the study. These endemic avian species not only emphasize the ecological importance of the reforested areas but also serve as bio indicators for habitat restoration success (Chowfin and Leslie, 2021).

Some research shows that the creation of new agricultural lands makes it imperative to take swift action in order to prevent the disappearance of corridor elements between fragments. The establishment of newly connected corridors may have the potential to revitalize pollinator communities and restore essential pollination services. (Kormann *et al.*, 2016). Perera *et al.* (2017) show that the potential value of wooded forest-home garden interfaces and forest stands restored with exotic pines in supporting native forest bird assemblages, especially in fragmented and isolated forest patches in the lowland rainforest in Sri Lanka. Further, some songbird species' territories were shrinking due to habitat fragmentation (Dilhari and Wickramasinghe, 2016).

These success stories illustrate the power of habitat restoration in turning the tide for avian species on the brink of extinction. By creating suitable habitats and addressing the threats they face, restoration efforts breathe new life into avifauna populations, underscoring the importance of continued conservation endeavors.

4. Conclusion

In conclusion, habitat restoration emerges as a powerful tool in rejuvenating avifauna populations and revitalizing our ecosystems. Throughout this journey, we have witnessed the profound impact of restoration efforts on our feathered friends.

First and foremost, habitat restoration has sparked recovery in bird populations, offering hope for the revival of avian biodiversity. As restored habitats provide abundant resources and suitable nesting sites, bird communities thrive, and the once-depleted skies resound with the joyful chorus of songbirds, waterfowl, and migratory species.

Moreover, restoration has demonstrated its ability to bolster nesting success for avian species. By creating safe and nurturing environments, we support avian parents in their vital task of raising the next generation, laying the foundation for sustainable populations that endure for generations to come.

Beyond the direct benefits to bird populations, habitat restoration breathes life into critical ecological processes. As avifauna engage in pollination, herbivory, seed dispersal, and decomposition, they orchestrate the symphony of life, ensuring the delicate balance of our ecosystems. The web of life begins to mend, as restored habitats pulsate with the rhythm of nature's interconnectedness.

However, the journey towards avian restoration does not end here. Ongoing research remains essential to refine our understanding of the intricate interactions between avifauna and restored ecosystems. By monitoring the responses of bird species and adapting management practices accordingly, we can optimize restoration efforts to yield even greater results.

As we embark on this endeavor, we call upon all nature enthusiasts, conservationists, and communities to take action. Supporting and participating in habitat restoration initiatives becomes an act of stewardship and a celebration of life's diversity. By volunteering, contributing, or simply raising awareness, we can each play a part in protecting avian biodiversity and preserving the symphony of nature's song for generations to come.

Let us unite in the spirit of restoration, knowing that every step we take towards reviving avian populations is a step towards restoring the harmony of our planet. Together, we can rewrite the narrative of biodiversity loss and compose a new melody where avifauna flourishes alongside us, forever etching their vibrant colors and graceful flights in the tapestry of our shared history.

References

Alsila, T., Elo, M., Hakkari, T., and Kotiaho, J.S. (2020). Effects of habitat restoration on peatland bird communities. *Restoration Ecology* :29;1 <https://doi.org/10.1111/rec.13304>

- Balvanera, P., C. Kremen, and M. Martinez-Ramos. (2005). Applying community structure analysis to ecosystem function: examples from pollination and carbon storage. *Ecological Applications* 15:360-375.
- Bollen, A., L. Van Elsacker, and J.U. Ganzhorn. (2004). Relations between fruits and disperser assemblages in a Malagasy littoral forest: a community-level approach. *Ecology* 20:599-612.
- Botes, A., M.A. McGeoch, and B. J. van Rensburg. (2006). Elephant and human-induced changes to dung beetle (Coleoptera: Scarabaeidae) assemblages in the Maputaland Centre of Endemism. *Biological Conservation* 130:573-583.
- Brawn, J. D. (2006). Effects of restoring oak savannas on bird communities and populations. *Conservation Biology* 20:460-469.
- Chapman, S. K., Hart, S. C., Cobb, N. S., Whitham, T. G., and Koch, G. W. (2003). Insect herbivory increases litter quality and decomposition: an extension of the acceleration hypothesis. *Ecology* 84:2867-2876.
- Chowfin, S.M and Leslie, A.J. (2021). Using birds as bioindicators of forest restoration progress: A preliminary study. *Trees, Forests and People* 3:100048
- Coley, P. D., and Barone, J. A. (1996). Herbivory and plant defenses in tropical forests. *Annual Review of Ecology and Systematics* 27:305-335.
- Dilhari, W. M. M. and Sriyani Wickramasinghe (2016). Fragmentation effects on territory quality of *Copsychus malabaricus* in the western part of the Mihintale sanctuary Sri Lanka, *Wildlanka*. 3 (3), 84-89
- Fink, R. D., Lindell, C. A., Morrison, E. B., Zahawi, R. A., and Holl, K. D. (2009). Patch size and tree species influence the number and duration of bird visits in forest restoration plots in southern Costa Rica. *Restoration Ecology* 17:479-486.
- Fletcher, R. J., and Koford, R. R. (2003). Changes in breeding bird populations with habitat restoration in northern Iowa. *American Midland Naturalist* 150:83-94.
- Fletcher, R. J., Koford, R. R., and Seaman, D. A. (2006). Critical demographic parameters for declining songbirds breeding in restored grasslands. *Journal of Wildlife Management* 70:145-157.
- Flesch A.D, Esquer A. (2020). Impacts of Riparian Restoration on Vegetation and Avifauna on Private and Communal Lands in Northwest Mexico and Implications for Future Efforts. *Air, Soil and Water Research*. 2020;13. doi:10.1177/1178622120938060
<https://doi.org/10.1177/1178622120938060>
- Gaines, W. L., Haggard, M., Lehmkuhl, J. F., Lyons, A. L., and Harrod, R. J. (2007). Short-term response of land birds to Ponderosa Pine restoration. *Restoration Ecology* 15:670-678.
- Gardali, T., Holmes, A. L., Small, S. L., Nur, N., Geupel, G. R., and Golet, G. H. (2006). Abundance patterns of land birds in restored and remnant riparian forests on the
- Higuchi H, Shibaev Y, Minton J, Ozaki K, Surmach S, Fujita G, Momose K, Momose Y, Ueta M, Andronov V, Mita N, Kanai Y. 1998. Satellite tracking of the migration of the red-crowned crane *Grus japonensis*. *Ecol Res*, 13:273-282
- Johnson, T. N., Applegate, R. D., Hoover, D. E., Gipson, P. S., and Sandercock, B. K. (2009). Evaluating avian community dynamics in restored riparian habitats with mark-recapture models. *Wilson Journal of Ornithology* 121:22-40.
- Johnson, T. N., and Sandercock, B. K. (2010). Restoring tallgrass prairie and grassland bird populations in tall fescue pastures with winter grazing. *Rangeland Ecology and Management* 63:679-688.
- Kormann U, Scherber C, Tschardtke T, Klein N, Larbig M, Valente JJ, Hadley AS, Betts MG. (2016). Corridors restore animal-mediated pollination in fragmented tropical forest landscapes. *Proc. R. Soc. B* 283: 20152347.
<http://dx.doi.org/10.1098/rspb.2015.2347>

- Macias, A. C. I. (2009). Effects of Habitat Fragmentation on the Distribution and Movement of Tropical Forest Birds. Open Access Dissertations:481.
- Majer, J. D. (2009). Animals in the restoration process: processing the trends. *Restoration Ecology* 17:315-319.
- Masters, R. E., Wilson, C. W., Cram, D. S., Bukenhofer, G. A., and Lochmiller, R. L. (2000). Influence of ecosystem restoration for Red-cockaded Woodpeckers on breeding bird and small mammal communities. In W. M. Ford, K. R. Russell, & C. E. Moorman (Eds.), *The Role of Fire in Non-game Wildlife Management and Community Restoration: Traditional Uses and New Directions* (pp. 73-90). Proceedings of a Special Workshop. USDA Forest Service Northeastern Research Station.
- Momose, S. and Momose, K.(2019). Species review:Red-crowned crane (*Grus japonensis*). IUCN SSC Crane Specialist Group – Crane Conservation Strategy
- Ortega-Alvarez, R., and Lindig-Cisneros, R. (2012). Feathering the scene: the effects of ecological restoration on birds and the role birds play in evaluating restoration outcomes. *Ecological Restoration* 30(2):116-127. doi:10.1353/ecr.2012.0038
- Ortega-Álvarez, R., Lindig-Cisneros, R., MacGregor-Forsb, I., Renton, K., and Schondube, J. E. (2013). Avian community responses to restoration efforts in a complex volcanic
- Perera, P., Wijesinghe, S., Dayawansa, N., Marasinghe, S., & Wickramarachchi, C. (2017). Response of tropical birds to habitat modifications in fragmented forest patches: A case from a tropical lowland rainforest in south-west Sri Lanka. *Community Ecology*, 18(2), 175–183. <https://www.jstor.org/stable/90017721>
- Ribeiro, M.C., Metzger, P.J., Martensen, A.C., Ponzoni, F.J. and Hirota, M.M. (2009). The Brazilian Atlantic Forest: How much is left, and how is the remaining forest distributed? Implications for conservation. *Biological Conservation* : 142(6), 1141-1153 <https://doi.org/10.1016/j.biocon.2009.02.021>
- Seigel, A. C., Hatfield, C., and Hattman, J. M. (2005). Avian responses to restoration of urban tidal marshes in the Hackensack Meadowlands, New Jersey. *Urban Ecosystems* 3:87-116.
- Sekercioglu, C. H. (2006). Increasing awareness of avian ecological function. *Trends in Ecology and Evolution* 21:464-471. SER. (2004).
- Small, S. L., Thompson, F. R., Geupel, G. R., and Faaborg, J. (2007). Spotted Towhee population dynamics in a riparian restoration context. *The Condor* 109:721-733.
- Twedt, D. J., Wilson, R. R., Henne-Kerr, J. L., and Grosshuesch, D. A. (2002). Avian response to bottomland hardwood reforestation: the first 10 years. *Restoration Ecology* 10:645-655.
- Wunderle, J. M. (1997). The role of animal seed dispersal in accelerating native forest regeneration of degraded tropical lands. *Forest Ecology and Management* 99:223-235.
- Young, T. P. (2000). Restoration ecology and conservation biology. *Biological Conservation* 92:73-83.