

study, it is concluded that this industry provides vast opportunity for further exploiting labour force in respect to production of beverage, sugar, alcohol, fibre, fuel wood, timber and raw materials for handicrafts. As well as this study clearly stated that implementing innovative technologies with product diversification and opening new market channels are the necessary pre requisite of this industry in future.

010

Effect of free range poultry system on land use efficiency and floral diversity in Rubber plantations

B M D C Balasooriya¹, V H L Rodrigo¹, S M M Iqbal¹ and D V S de S Gamage²,

¹Rubber Research Institute of Sri Lanka, ²Veterinary Research Institute, Sri Lanka

Plantation crops like rubber were traditionally grown in monocultures to avoid any risk on their productivity. However, with the smallholder farmers being involved in rubber cultivation, the idea of farming system evolved with the maximizing the overall land productivity and diversifying the income sources in order to meet the need of the resource poor. At present, rubber based intercropping systems with other economically beneficial crops are recommended and being practised to some extent particularly in smallholdings. However, no successful attempts were taken on livestock integration to the rubber system. With the understanding of practical limitations in incorporating ruminants to the system, the present study was commenced as a preliminary investigation to assess the effectiveness of free range poultry with respect to productivity and its effects on floral diversity under rubber. Based on the experience in coconut based poultry systems in Sri Lanka, a strain developed as a backcross of the hybrid between CPRS (Central Poultry Research Station) Brown and indigenous with indigenous, was used in this study expecting desired characteristics of both types, i.e., high level of egg production and adaptability to the environment. The trial began with 30 birds in a mature rubber clearing of the Rubber Research Institute of Sri Lanka. Information on egg production, the effect poultry on weed growth and its diversity were recorded together with visual observations on birds' behavior.

Average egg production was 8 eggs per bird per month. This value has dropped to 3 when the majority of birds were in moulting. However, in some months, it has gone up to 14. Some birds were killed by predators and even unsuccessful attacks affected the egg production. Birds showed poor brooding characters with that attempts taken to hatch eggs were failed. Poultry feeds had to be supplied (50g layer ration per bird) to maintain the continuous egg production. Birds always used to feed on the area close human dwelling. Dominance of the floral species under rubber was changed with the incorporation of poultry. Although *Adiantum latifolium* was dominant throughout the study, Summed Dominance Ratio (SDR) of *Paspalum conjugatum* and *Syngonium podophyllum* declined with the poultry integration. Measures to be taken to improve the poultry system were also identified.

011

Rubber latex production in *Hevea brasiliensis* with high density planting

T U K Silva¹, V H L Rodrigo¹, S M C U P Subasinghe²

¹Rubber Research Institute of Sri Lanka,

²Department of Forestry and Environmental Science, University of Sri Jayewardenepura, Sri Lanka.

Despite short-term fluctuation of rubber prices, the demand of natural rubber has increased continuously with the increase in population and living standards of the human being. Nevertheless, urbanization results in the fast depletion of forests as well as the land area under rubber. Therefore, in order to meet the continuous demand on latex, the productivity of rubber plantations should be increased. While producing high yielding clones for improved yield per tree which is a long-term process in perennial crops, planting density could be adjusted to obtain high productivity in rubber plantations. The present level of planting density of rubber in Sri Lanka has been decided on the experiments conducted with the genotypes which are not in common use at the moment. Also, the optimum density should vary with different socio-economic conditions. Therefore, the present study was aimed to identify the suitable planting density for the recently developed and commonly used genotypes of

rubber. Specifically, this paper is focused to assess the overall productivity of rubber with respect to latex yield.

The experiment was set up in Rathnapura district of Sri Lanka in 1992. Rubber was planted in three high densities, i.e., 600, 700, 800 trees per hectare, with the presently recommended level of 500 trees per hectare. Also, 3 genotypes (clones) i.e., RRIC 100, RRIC 110 and RRIC 121 were incorporated with the statistical design of split plot where the planting densities were laid as the main plots whilst clones were in the sub plots. Growth and yield parameters in terms of girth, bark thickness (BT), and the incidences of tapping panel dryness (TPD) were assessed yearly and the latex volume (LV), % dry rubber content (DRC), number of trees in tapping (TIT) were assessed on daily basis up to 2004.

Irrespective of the clone used, mean yield per tree per tapping decreased with increase in planting density. However, it was *vice versa* in the case of yield per hectare (YPH) due to the increase in TIT with increase in planting density. The percentage of trees with TPD was not significantly affected by the planting density. The clone RRIC 110 was infected with *Corynespora* leaf disease hence poor performance was shown in all densities. The trend of increasing YPH was similar in both other clones, i.e. RRIC 100 and RRIC 121. However, the rate of increase in TIT showed a decline of 700 trees per hectare resulting in a lower YPH at this level than expected. Although latex productivity could generally be increased with increase in planting density, overall economic profitability of the system will also depend on cost of production, amount and value of timber and carbon produced. Therefore, study warrants further investigations on above issues before making any changes to the presently recommended planting density.

012

Assessment on timber and carbon in rubber plantations with special reference to the wet zone of Sri Lanka

E S Munasinghe¹, V H L Rodrigo¹, P K W Karunathilaka¹ and U A D P Gunawardena²

¹ Rubber Research Institute of Sri Lanka.

² Department of Forestry and Environmental Science, University of Sri Jayewardenepura, Sri Lanka

Rubber (*Hevea brasiliensis*) has traditionally been cultivated for the latex extraction; however, its importance in other uses, particularly producing quality timber and sequestering atmospheric carbon as a permanent sink, is also often highlighted. The amount of timber produced and carbon sequestered in rubber trees has been assessed in isolations. Those values would differ under different growth conditions and to date, no simple protocol is available to quantify the amount of timber and carbon in rubber plantations. Therefore, the study reported here was aimed to develop simple growth models to assess the timber production and carbon fixing capability of rubber plantations in Sri Lanka.

Initially, a growth function was developed to assess the girth development with respect to age and thereafter another three functions to quantify the amount of timber, biomass and carbon in the rubber tree based on girth diameter. Also, wood density variation with age of the tree was modeled to determine the biomass in timber under different age categories. The assessment on the available carbon was based on the carbon content in unit biomass and the total amount of biomass in the tree. Growth data required for the girth development function were gathered from secondary sources and girth measurements made on existing rubber clearings. Destructive sampling was conducted to assess the timber, biomass and wood density.

Based on above models, an average rubber tree at 30 years achieves a girth of 88.64 cm and produces 0.656 m³ of timber and 594.46 kg of biomass. The amount of atmospheric carbon fixed in timber at this age was estimated as 193.7 kg per tree and 45.86 MT per hectare. However, total amount of organic carbon fixed in above ground components was 220.8 kg per tree and 52.27 MT per hectare. The models of this study were developed under general conditions in the wet zone, hence should be validated for drier regions of the country before any wide scale adoption.